


(Fl-control, Fl-anti-V alpha 3, Fl-anti-TCR alpha/beta, Fl-anti-Vbeta 8) Immunofluorescence of MCA-26 tumor cells






Fig. 6 a) (inset) A W/Fu rat bearing SMT-2A shows massive, generalized lymph node metastasis. b) SMT-2A cells immuno-cytochemically stained with mouse monoclonal antibody specific to the rat CD8 T-cells. The secondary antibody was labeled with FITC.

ME = LYMPH NODE HETASTASIS

P = PRIMARY SMT-ZA TUHOR

the term that the term to the






Fig. 7. Growth characteristics of SW480E and SW480R colon tumor cells in athymic nude mice







Days after subcutaneous injection of 2X10<sup>6</sup> SW480E tumor cells

^ = IP injection of anti-AsGM1


nation, nations


**F**<sub>g</sub>.9





ĘII






anti - C $\!\beta$ 

anti - Vβ3

anti - IgG

SW480










SW620










F1915



D

<u>|</u> \_5

أإزا

H

IJ

1

II

21

11

31

71

241

51

Fg. 17

A -72 SYK -70 SYKB

F<sub>y</sub>. 18



Effect of anti-V beta 3 on proliferation of SW620 cells in serum free media







Fig. 20
Effect of IL-16 on proliferation of SW620 cells in serum free media






STAGE-I REAST CANCER CELLS IN 75 YEAR D WOMAN CONTAINING MANY (88.7%) CD3 (T cell receptor) POSITIVE TUMOR CELLS.



FLUORESCENCE PHOTOMICROGRAPHS OF CD3+EMA+ HUMAN MAMMARY CARCINOMA CELLS.



PE=Phytoerythrin (red dye); FITC=Fluorescene isothiocyanate (green dye). EMA=Epithelial membrane antigen (indicates these cells are breast cancer cells).